

## Tropical Seagrass as a Bioindicator of Metal Accumulation (Rumput Laut Tropika sebagai Penunjuk Biologi Pengumpulan Logam)

FARIDAHANIM AHMAD, SHAMILA AZMAN\*, MOHD ISMID MOHD SAID & LAVANIA-BALOO

### ABSTRACT

*Seven species of tropical seagrass found at seagrass bed located in Johor, Malaysia were analysed for As, Cu and Cd accumulation. The species were identified as Enhalus acoroides, Halophila minor, Halophila spinulosa, Halophila ovalis, Thalassia hemprichii, Halodule uninervis and Cymodocea serrulata. Seagrass plant is rapidly becoming one of the methods to determine the overall health condition of aquatic environment. Each seagrass samples were collected and divided into three parts i.e roots, rhizomes and leaves. Samples were grinded, digested and the correlation between each part was analysed using SPSS version 16. Each part of seagrass tissues have the ability to assimilate metals for example the concentration of As, Cu and Cd in tropical seagrass were in the range of 5-48, 6-60 and 10-69 µg/gDW, respectively. Halophila minor and Halophila ovalis indicates positive correlations to translocate metals (As, Cu and Cd) in plants parts (leaves-rhizomes, rhizomes-roots and roots-leaves). Seagrass can accumulate metals depending on pollution that occur, seasonal variation and internal capabilities to translocate metals. The seagrass species especially Halophila ovalis and Halophila minor can act as bioindicator for metal pollution.*

*Keywords: Bioindicator; metal contamination; seagrass*

### ABSTRAK

*Tujuh spesies rumput laut tropika yang dijumpai di beting rumput laut yang terletak di Johor, Malaysia telah dianalisis untuk pengumpulan logam As, Cu dan Cd. Spesies yang dikenal pasti adalah Enhalus acoroides, Halophila minor, Halophila spinulosa, Halophila ovalis, Thalassia hemprichii, Halodule uninervis dan Cymodocea serrulata. Rumput laut boleh dijadikan sebagai salah satu kaedah untuk menentukan keadaan kesihatan keseluruhan persekitaran akuatik. Setiap sampel rumput laut dikumpulkan dan dibahagikan kepada tiga bahagian iaitu akar, rizom dan daun. Sampel dikisar, dicerna dan korelasi antara setiap bahagian telah dianalisis menggunakan perisian SPSS versi 16. Setiap bahagian tisu rumput laut mempunyai keupayaan untuk mengasimilasikan logam contohnya keputusan menunjukkan kepekatan As berada dalam julat 5-48 µg/gDW, Cu dalam julat 6-60 µg/gDW dan Cd dalam julat 10-69 µg/gDW. Halophila minor dan Halophila ovalis menunjukkan korelasi positif untuk memindahkan logam (As, Cu dan Cd) pada tumbuhan (daun-rizom, akar rizom dan akar-daun). Rumput laut tropika boleh mengumpul logam bergantung kepada pencemaran yang berlaku, perbezaan musim dan keupayaan dalaman rumput laut untuk memindahkan logam. Spesies rumput laut terutama Halophila ovalis dan Halophila minor boleh dijadikan sebagai petunjuk biologi.*

*Kata kunci: Pencemaran logam; penunjuk biologi; rumput laut*

### INTRODUCTION

Estuaries are exposed to anthropogenic contaminants including complex mixtures of heavy metals from industrial, agricultural and domestic waste directly via rivers or through atmospheric deposition (Emelogu et al. 2013; Gillet et al. 2008; Lafabrie et al. 2007; Villate et al. 2013). Human activities such as coastal land disturbance, motor boating and dredging increases pollutants (Burkholder et al. 2007; Lewis et al. 2007). One way to measure the degree of pollution is by using seagrass as a bioindicator for metal contamination (Ferrat et al. 2003; Prange & Dennison 2000; Schlacher-Hoenlinger & Schlacher 1998). Seagrass species are rooted angiosperms marine plants that are widely distributed in a large area known as seagrass bed (Bastyan & Cambridge 2008;

Kirkman 1990; McKenzie et al. 2001; Vermaat et al. 2004). Seagrass bed plays an ecological role in coastal ecosystems as the most productive plant communities that support high biodiversity and food for a variety of marine organisms (Duarte & Chiscano 1999; Eklöf et al. 2008; Jackson et al. 2006; Lee et al. 2007; Waycott et al. 2005). Seagrass plant is rapidly becoming one of the method to determine the overall health and condition of aquatic environment (Bortone & Turpin 2000; Costantini et al. 1991; Macinnis-Ng & Ralph 2004). Seagrass have a remarkable metal bioaccumulation capacity since it interacts directly with both water column and pore water through the leaves and roots as ionic uptake hence it can reflect the overall health of coastal water (Llagostera et al. 2011). Seagrasses are protected under International Union

for the Conservation of Nature Red list of Threatened Species and the biggest diversity of tropical seagrass was identified at Indo-Pacific region (Short et al. 2011, 2001). In addition, the seagrass bed serves as a 'nursery habitat' for the most productive plant communities that support food and shelter for a diverse biodiversity and vulnerable marine organisms (Duarte & Chiscano 1999; Heck & Valentine 2006, Jackson et al. 2006; Lee et al. 2007; Waycott et al. 2005). However, the continuously increasing pollution will cause the seagrasses to die-off once pollution reaches lethal levels and this intertidal habitat will be lost for good as there is little opportunity for the habitat itself to migrate (Hadley 2009).

Currently data recorded for metal accumulation in tropical seagrass in Southeast Asia is very limited (Ooi et al. 2011). Therefore, the aim of this study was to determine the capability of 7 tropical seagrass species (*Enhalus acoroides*, *Halophila minor*, *Halophila spinulosa*, *Halophila ovalis*, *Thalassia hemprichii*, *Halodule uninervis* and *Cymodocea serrulata*) at Pulau River Estuary to accumulate As, Cu and Cd. The seagrass bed at Pulau River Estuary is the largest in Malaysia with an approximate size area of 3.15 km<sup>2</sup>. The estuary is listed as a Ramsar Site and receives water from the Straits of Malacca and nearby rivers (Rusli 2012). It is located nearby Malaysia - Singapore Second Link, Port of Tanjung Pelepas and Tanjung Bin Power Plant.

## MATERIALS AND METHODS

### SAMPLING

Seagrass samples were collected at Pulau River Estuary seagrass bed from July, 2011 to March, 2012. The samples complete with roots, rhizomes and leaves were manually collected at every 20 m interval on the seagrass bed using transect method. The starting sampling point on seagrass bed is at 1° 20' 22.53" N, 103° 36' 16.13" E. Pulau seagrass bed is an intertidal zones in which during high tide the depth from surface of seawater to seagrass bed is approximately 3.5 m. Figure 1 shows the location, perimeter, upper view and lateral view of Pulau seagrass bed. The slope gradients were gradually increasing and it can be visually seen like a big patch of grassland during low tide. The gradient shows a gradual slope at every edge of the seagrass bed. The size of the seagrass bed is 1.8 km in length by 0.1 km width. The distance between seagrass bed to Port of Tanjung Pelepas, Tanjung Bin Power Plant and Bunker Terminal are 3.7, 5.6 and 5.5 km, respectively.

The samples collected were later rinsed with seawater to remove sediment at roots and rhizomes. Then, the samples were packed in clean plastic bags, sealed, placed in a cool box containing ice, then transported to the laboratory and frozen at -20°C prior to analysis (Sanchiz et al. 2000).




FIGURE 1. a) Location of the seagrass bed relative to the Port of Tanjung Pelepas, Tanjung Bin Power Plant and the Malaysia-Singapore Second link bridge, b) Pulau Estuary seagrass bed: the yellow-dotted irregular oval is the perimeter of the seagrass bed (goggle earth), c) lateral view of the seagrass bed on site during low tide, the gradient is gradually slope at the edge of seagrass bed and d) upper view of the seagrass bed (google earth)

## SAMPLE PROCESSING AND DATA ANALYSIS

Each seagrass sample was collected and mixed at different locations on the seagrass bed. The sample was sorted into leave, rhizome and root. The seagrass samples were collected in a set of 12 specimens. Samples were dried at room temperature until it reaches a constant weight and grinded using agate mortar into homogenous powder (Llagostera et al. 2011). 0.3 g of the sample was later placed in Pyrex tubes with Teflon closure and added with 5 mL  $\text{HNO}_3$ , incubated for 1 h in a water bath at  $100^\circ\text{C}$ , then cooled, filtered with  $0.2\ \mu\text{m}$  nylon membrane and diluted in 15 mL deionised water. Metal analysis was carried out using Perkin Elmer Atomic Absorption Spectrophotometer Model AAnalyst 400.

The Pearson's correlation coefficients of metals concentration between roots, rhizomes and leaves tissues were analysed using SPSS version 16 software. For easy references and recognition the seagrasses were given code names as shown in Table 1.

## RESULTS AND DISCUSSION

Figure 2 shows the concentration of As, Cu and Cd in different tissues of the 7 tropical seagrass. The results were expressed as  $\mu\text{g}$  of metal per gDW of analysed tissue where DW refers to dry weight. All seagrass species were able to accumulate As, Cu and Cd. The concentration of As, Cu and Cd in different tissues of the 7 tropical

TABLE 1. Code names of seagrasses

| Code name | Species of seagrass         |
|-----------|-----------------------------|
| Ea        | <i>Enhalus acoroides</i>    |
| Hm        | <i>Halophila minor</i>      |
| Hs        | <i>Halophila spinulosa</i>  |
| Ho        | <i>Halophila ovalis</i>     |
| Th        | <i>Thalassia hemprichii</i> |
| Hu        | <i>Halodule uninervis</i>   |
| Cs        | <i>Cymodocea serrulata</i>  |

seagrass were in the range of 5-48, 6-60 and 10-69  $\mu\text{g/gDW}$ , respectively.

At Pulau River Estuary, seagrass tissue can assimilate higher Cd content rather than Cu and As. Cd sources may come from Port of Tanjung Pelepas, Bunker Terminal and the Straits of Malacca which can cause the reclamation work at the estuary and increasing shipping activities. The Cd discharged and flow into the seawater will then uptake by the seagrass. Marine plants that accumulate Cd from seawater is considered as toxic at higher concentration (Ripperger et al. 2007).

Cu can also be accumulated in all the seven species of tropical seagrass. Bioavailability and toxicity of Cu to aquatic organisms depends on the total concentration at the estuary. Elevated Cu concentration in estuarine environments may result from the industrial and municipal waste, urban runoff, rivers and shorelines




FIGURE 2. Concentration of As, Cu and Cd in seagrass tissues; all data express by  $\mu\text{g/gDW}^{-1}$ ; Mean and standard error are shown

erosion related to human activities from Pulai River. Cu is an essential component of many proteins, in excess it can cause toxicity and lead to damaging effect in marine plant tissue particularly to mechanisms involved in photosynthesis (Grant et al. 2003). In addition, previous study reported that Cu is toxic to many phytoplankton and zooplankton at high concentration (Vance et al. 2008).

Tropical seagrasses are able to assimilate As from seawater. The source of As could come from the biggest coal-fired in Southeast Asia known as Tanjung Bin Power Plant. As is toxic and has no biological role (Rensing & Rosen 2009). As appears to be elevated in marine biota because of its ability to accumulate As from seawater. Macrophytes transform arsenate into non-volatile methylated As compounds such as methanearsonic and dimethylarsinic acids, methylation of As (unlike methylation of Hg) greatly reduced toxicity and is known as detoxification process (Eisler 2000).

Undesired metal contaminates (As, Cu and Cd) were trapped on seagrass bed due to the increasing gradient surface of sediment. All seagrass tissues could reasonably have a bioindicator role in reducing metals contamination that comes from seawater and sediment, therefore partially can reduce the effect of pollution on Pulai River Estuary. The pollutants are believed to come from nearby development, land use and shipping oil spills from the Malacca and Singapore Straits. The seagrass bed's ability to trap metal contaminants is evident that it is functioning well, as stated by Short and Coles (2001) where seagrass bed acts as a buffer zone. Aside from that, the seagrass bed cleanses seawater by absorbing dissolved metals and is a bioindicator that reflects the overall health of the aquatic ecosystem (Costantini et al. 1991; Eklöf et al. 2008; Kaldy 2006; Llagostera et al. 2011; Macinnis-Ng and Ralph 2004).

Table 2 shows the concentration of As, Cu and Cd from the previous studies for temperate and tropical seagrasses. In this paper, Cu and Cd accumulates higher concentration in *Cymodocea serrulata*, *Thalassia hemprichii*, *Enhalus acoroides*, *Halodule uninervis* and *Halophila ovalis* than reported in Flores Sea, Indonesia (Nienhuis 1986). The different value of accumulation is due to the different metals rate at surrounding area. Flores Sea is situated without industry and untouched isolated areas whereas Pulai River Estuary seagrass bed is surrounded by on-going development between Malaysia and Singapore. Metals concentrations in tropical seagrass are basically depend on the metals presence either the area is polluted or not.

The Cd concentrations reported by Kilminster (2013) were lower compared with this study. The concentrations were different because of seasonal variation. At Pulai River Estuary, the weather condition is always warm and the range of seawater temperature is  $27.0 \pm 3.0^\circ\text{C}$  whereas Kilminster (2013) collected his samples in winter. As stated by Eisler (2000) Cd uptake will increase with

the increasing seawater temperature in a range of 5 to  $25^\circ\text{C}$ . Cd concentration in this study is similar to studies reported by Malea (1994) and Malea et al. (1994) where the seawater temperature is in the range of  $11\text{-}24^\circ\text{C}$ .

The concentrations of As and Cu in the seagrass tissues are also similar to the results reported by previous studies (Eisler 2000; Kilminster 2013; Malea 1994; Malea et al. 1994; Schlacher-Hoenlinger & Schlacher 1998). Generally, bioavailability and toxicity of marine plants that accumulates As, Cu and Cd from seawater and sediment are dependent on its total metal concentrations in the estuary and seasonal variation (Eisler 2000; Malea et al. 1994). Seagrass tissues are considered toxic at higher metal concentration because it tends to accumulate metals from seawater and sediment to reduce metal contamination from the surroundings (Macinnis-Ng & Ralph 2002; Ripperger et al. 2007).

Table 3 shows the correlations of As in roots-rhizomes, rhizomes-leaves and leaves-roots, where there are significant correlations between different seagrass tissues for *Enhalus acoroides*, *Halophila minor*, *Halophila ovalis* and *Cymodocea serrulata*. There are significant correlations for accumulation of Cu between roots-rhizomes, rhizomes-leaves and leaves-roots in *Halophila minor*, *Halophila ovalis*, *Halodule uninervis* and *Cymodocea serrulata* (Table 4). As for Cd accumulation (Table 5), only *Halophila minor*, *Halophila ovalis* and *Cymodocea serrulata* were found to have significant correlations between roots-rhizomes, rhizomes-leaves and leaves-roots. Based on the correlation analysis, it can be summarized that in *Halophila minor*, *Halophila ovalis* and *Cymodocea serrulata* there are positive correlations between roots, rhizomes and leaves in metals (As, Cu and Cd). This indicate that *Halophila minor*, *Halophila ovalis* and *Cymodocea serrulata* have effective active uptake kinetics and internal tissues to translocate As, Cu and Cd to the entire plant system.

The correlation data (As, Cu and Cd) for leaves, rhizomes and roots are different because metal accumulation by seagrass are governed by multitudes of factors such as the relative metal availability for seagrass uptake, plant physiology including leaves may differ uptake kinetics and passive absorption from roots and internal translocation among plant parts, be it passively or actively (Llagostera et al. 2011). Active uptake is a movement and transported biochemical from lower concentration to higher concentration that requires energy known as uphill process whereas passive uptake known as downhill process are driven by mass flow (Zhan et al. 2010). There are two phases for absorbing metals for terrestrial plant, the first phase is the metals diffuse from the surrounding into the root; in the second phase metals gradually distributes and accumulates in plant tissues (Kvesitadze et al. 2009). In this study, seagrass tissues (roots, rhizomes and leaves) may absorb the metals simultaneously because the seagrass were submerged in seawater.

TABLE 2. Concentration of As, Cu and Cd from temperate and tropical seagrass species in  $\mu\text{g DW}^{-1}$ 

| Seagrass   | Location                  |               | Metals concentration |           |           |                     | References                            |
|--|---------------------------|---------------|----------------------|-----------|-----------|---------------------|---------------------------------------|
|  |                           |               | Arsenic              | Copper    | Cadmium   |                     |                                       |
| Temperate  | <i>Posidonia oceanica</i> | Whole plant   | Mediterranean        | -         | -         | 5.38±0.14           | Lafabrie et al. 2007                  |
|  |                           | Whole plant   | Mediterranean        | -         | -         | 3.39±0.12           | Lafabrie et al. 2007                  |
|  |                           | Whole plant   | Mediterranean        | -         | -         | 1.70±0.06           | Lafabrie et al. 2007                  |
|  |                           | Whole plant   | Italy                | -         | 14.0±17.0 | 0.9-1.3             | Schlacher-Hoenlinger & Schlacher 1998 |
| <i>Halophila stipitata</i>                                   | Whole plant               | Greece        | -                    | 2.0-82.7  | 1.3-93.7  | Malea 1994          |                                       |
|  | Root and rhizome          | Greece        | -                    | 1.9-79.7  | 0.88-51.7 | Malea 1994          |                                       |
|  | Leaf                      | Greece        | -                    | 1.9-81.3  | 2.5-85.7  | Malea 1994          |                                       |
| <i>Posidonia oceanica</i>                                    | Leaf                      | Mediterranean | -                    | -         | 2.1-25.0  | Sanchiz et al. 2000 |                                       |
|  | Rhizome                   | Mediterranean | -                    | -         | 0.6-2.0   | Sanchiz et al. 2000 |                                       |
|  | Root                      | Mediterranean | -                    | -         | 0.9-2.2   | Sanchiz et al. 2000 |                                       |
|  | Whole plant               | Greece        | -                    | 2.8-148.0 | 2.7-44.0  | Malea et al. 1994   |                                       |
| Tropical   |                           |               |                      |           |           |                     |                                       |
| <i>Cymodocea rotundata</i>                                   | Leaf                      | Indonesia     | -                    | 6.6±1.6   | 0.49±0.24 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 4.5±1.6   | 0.40±0.02 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 4.5±5.0   | 0.22±0.03 | Nieuhuis 1986       |                                       |
| <i>Cymodocea serrulata</i>                                   | Leaf                      | Indonesia     | -                    | 5.8±2.2   | 0.68±0.15 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 3.1±1.0   | 0.27±0.08 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 3.1±1.8   | 0.17±0.03 | Nieuhuis 1986       |                                       |
| <i>Thalassia hemprichii</i>                                  | Leaf                      | Indonesia     | -                    | 7.0±2.3   | 0.57±0.33 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 5.0±2.0   | 0.42±0.28 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 2.9±1.2   | 0.24±0.19 | Nieuhuis 1986       |                                       |
| <i>Syringodium isoetifolium</i>                              | Leaf                      | Indonesia     | -                    | 3.9±1.8   | 0.16±0.63 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 4.1±1.7   | 0.21±0.08 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 6.8±8.1   | 0.16±0.05 | Nieuhuis 1986       |                                       |
| <i>Enhalus acoroides</i>                                     | Leaf                      | Indonesia     | -                    | 4.4±2.3   | 0.36±0.19 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 3.7±2.3   | 0.16±0.12 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 2.6±2.3   | 0.12±0.07 | Nieuhuis 1986       |                                       |
| <i>Halodule uninervis</i>                                    | Leaf                      | Indonesia     | -                    | 5.6±2.4   | 0.60±0.19 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 3.4±3.0   | 0.24±0.07 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 2.7±0.4   | 0.25±0.07 | Nieuhuis 1986       |                                       |
| <i>Halodule pinifolia</i><br><i>Thalassodendron ciliatum</i> | Whole plant               | Indonesia     | -                    | 6.7±5.2   | 0.36±0.21 | Nieuhuis 1986       |                                       |
|  | Leaf                      | Indonesia     | -                    | 6.0±0.9   | 1.54±0.56 | Nieuhuis 1986       |                                       |
|  | Shoot                     | Indonesia     | -                    | 3.3±0.9   | 0.98±0.20 | Nieuhuis 1986       |                                       |
|  | Rhizome and root          | Indonesia     | -                    | 3.4±0.7   | 0.73±0.39 | Nieuhuis 1986       |                                       |
| <i>Halophila ovalis</i>                                      | Whole plant               | Indonesia     | -                    | 17.3±15.1 | 0.41±0.12 | Nieuhuis 1986       |                                       |
|  | Leaf                      | Australia     | 5.4-8.5              | 6.6-42.2  | 0.18-0.53 | Kilminster 2013     |                                       |
|  | Rhizome                   | Australia     | 9.2-20.0             | 3.3-26.6  | 0.04-0.38 | Kilminster 2013     |                                       |
|  | Shoot                     | Australia     | 15.4-35.0            | 11.1-80.0 | 0.05-0.38 | Kilminster 2013     |                                       |

TABLE 3. Correlation analysis for As concentration in roots, rhizomes and leaves with r: correlation coefficient; p-value: significance level settled at 0.05 (2-tailed); n=12

|    | Roots-rhizomes |         | Rhizomes-leaves |         | Leaves-roots |         |
|----|----------------|---------|-----------------|---------|--------------|---------|
|    | r              | p-value | r               | p-value | r            | p-value |
| Ea | 0.92           | 0.00    | 0.98            | 0.00    | 0.98         | 0.00    |
| Hm | 0.99           | 0.00    | 1.00            | 0.00    | 1.00         | 0.00    |
| Hs | 0.90           | 0.01    | 0.55            | 0.20    | 0.86         | 0.01    |
| Ho | 0.98           | 0.00    | 1.00            | 0.00    | 0.98         | 0.00    |
| Th | 0.17           | 0.14    | 0.93            | 0.00    | 0.06         | 0.39    |
| Hu | 0.51           | 0.05    | 0.56            | 0.03    | 0.04         | 0.62    |
| Cs | 1.00           | 0.00    | 1.00            | 0.00    | 1.00         | 0.00    |

Table 4. Correlation analysis for Cu concentration in roots, rhizomes and leaves with r: correlation coefficient; p-value: significance level settled at 0.05 (2-tailed); n=12

|    | Roots-rhizomes |         | Rhizomes-leaves |         | Leaves-roots |         |
|----|----------------|---------|-----------------|---------|--------------|---------|
|    | r              | p-value | r               | p-value | r            | p-value |
| Ea | 0.67           | 0.10    | 0.72            | 0.07    | 0.99         | 0.00    |
| Hm | 0.98           | 0.00    | 1.00            | 0.00    | 0.97         | 0.00    |
| Hs | 0.02           | 0.96    | 0.82            | 0.02    | 0.88         | 0.00    |
| Ho | 1.00           | 0.00    | 0.86            | 0.01    | 0.87         | 0.01    |
| Th | 0.49           | 0.26    | 0.03            | 0.55    | 0.28         | 0.05    |
| Hu | 1.00           | 0.00    | 1.00            | 0.00    | 1.00         | 0.00    |
| Cs | 1.00           | 0.00    | 1.00            | 0.00    | 0.99         | 0.00    |

TABLE 5. Correlation analysis for Cd concentration in roots, rhizomes and leaves with r: correlation coefficient; p-value: significance level settled at 0.05 (2-tailed); n=12

|    | Roots-rhizomes |         | Rhizomes-leaves |         | Leaves-roots |         |
|----|----------------|---------|-----------------|---------|--------------|---------|
|    | r              | p-value | r               | p-value | r            | p-value |
| Ea | 0.94           | 0.01    | 0.16            | 0.20    | 0.71         | 0.00    |
| Hm | 0.99           | 0.00    | 1.00            | 0.00    | 0.99         | 0.00    |
| Hs | 0.40           | 0.03    | 0.58            | 0.23    | 0.12         | 0.27    |
| Ho | 0.99           | 0.00    | 1.00            | 0.00    | 1.00         | 0.00    |
| Th | 0.86           | 0.00    | 0.90            | 0.02    | 0.16         | 0.20    |
| Hu | 0.84           | 0.01    | 0.50            | 0.67    | 1.00         | 0.00    |
| Cs | 0.99           | 0.00    | 1.00            | 0.00    | 0.93         | 0.00    |

## CONCLUSION

In selecting seagrass species as a bioindicator, it should be taken into consideration that the highest accumulation of As is observed in *Halophila minor* (roots), the highest accumulation of Cu is in *Halophila minor* (rhizomes) and the highest accumulation of Cd is in *Halophila ovalis* (leaves). *Halophila minor* and *Halophila ovalis* also indicates a positive correlation to translocation of metals among plants parts, therefore *Halophila minor* and *Halophila ovalis* can be useful as bioindicators.

Two areas of research are identified as important consequences of this study. The various tissue structures of the seagrasses ought to be examined in order to explain their different absorption potentials, also the lethal concentration level for different seagrasses and for different metals should be further analysed.

## ACKNOWLEDGEMENTS

The authors wish to thank the Ministry of Education (MOE), Malaysia and Universiti Teknologi Malaysia for the financial support (UTM RUG Vot 03J65). We would also like to thank Professor Emeritus Gustaf Olsson for meaningful discussions.

## REFERENCES

- Bastyan, G.R. & Cambridge, M.L. 2008. Transplantation as a method for restoring the seagrass *Posidonia australis*. *Estuarine, Coastal and Shelf Science* 79: 289-299.
- Bortone, S.A. & Turpin, R.K. 2000. Tape grass life history metrics associated with environmental variables in a controlled estuary. In *Seagrass Monitoring, Ecology, Physiology, and Management*, edited by Bortone, S.A. Boca Raton: CRC Press LLC. pp. 65-79.

- Burkholder, J.M., Tomasko, D.A. & Touchette, B.W. 2007. Seagrasses and eutrophication. *Journal of Experimental Marine Biology and Ecology* 350: 46-72.
- Costantini, S., Giordano, R., Ciaralli, L. & Beccaloni, E. 1991. Mercury, cadmium and lead evaluation in *Posidonia oceanica* and *Codium tomentosum*. *Marine Pollution Bulletin* 22: 362-363.
- Duarte, C.M. & Chiscano, C.L. 1999. Seagrass biomass and production: A reassessment. *Aquatic Botany* 65: 159-174.
- Eisler, R. 2000. *Handbook of Chemical Risk Assessment Health Hazards to Human, Plants, and Animals*. Three volume set. Boca Raton: CRC Press.
- Eklöf, J.S., De La Torre-Castro, M., Gullström, M., Uku, J., Muthiga, N., Lyimo, T. & Bandeira, S.O. 2008. Sea urchin overgrazing of seagrasses: A review of current knowledge on causes, consequences, and management. *Estuarine, Coastal and Shelf Science* 79: 569-580.
- Emelogu, E.S., Pollard, P., Dymond, P., Robinson, C.D., Webster, L., Mckenzie, C., Dobson, J., Bresnan, E. & Moffat, C.F. 2013. Occurrence and potential combined toxicity of dissolved organic contaminants in the Forth estuary and Firth of Forth, Scotland assessed using passive samplers and an algal toxicity test. *Science of The Total Environment* 461-462: 230-239.
- Ferrat, L., Pergent-Martini, C. & Roméo, M. 2003. Assessment of the use of biomarkers in aquatic plants for the evaluation of environmental quality: Application to seagrasses. *Aquatic Toxicology* 65: 187-204.
- Gillet, P., Mouloud, M., Durou, C. & Deutsch, B. 2008. Response of *Nereis diversicolor* population (Polychaeta, Nereididae) to the pollution impact - Authie and Seine estuaries (France). *Estuarine, Coastal and Shelf Science* 76: 201-210.
- Grant, A.J., Graham, K., Frankland, S. & Hinde, R. 2003. Effect of copper on algal-host interactions in the symbiotic coral *Plesiastrea versipora*. *Plant Physiology and Biochemistry* 41: 383-390.
- Hadley, D. 2009. Land use and the coastal zone. *Land Use Policy* 26 (Supplement 1): S198-S203.
- Heck, J.K.L. & Valentine, J.F. 2006. Plant-herbivore interactions in seagrass meadows. *Journal of Experimental Marine Biology and Ecology* 330: 420-436.
- Jackson, E.L., Attrill, M.J., Rowden, A.A. & Jones, M.B. 2006. Seagrass complexity hierarchies: Influence on fish groups around the coast of Jersey (English Channel). *Journal of Experimental Marine Biology and Ecology* 330: 38-54.
- Kaldy, J.E. 2006. Carbon, nitrogen, phosphorus and heavy metal budgets: How large is the eelgrass (*Zostera marina* L.) sink in a temperate estuary? *Marine Pollution Bulletin* 52: 342-353.
- Kilminster, K. 2013. Trace element content of seagrasses in the Leschenault Estuary, Western Australia. *Marine Pollution Bulletin* 73: 381-388.
- Kirkman, H. 1990. Seagrass distribution and mapping. In *Seagrass Research Method*, edited by Phillips, R.C. & McRoy, C.P. France: United Nations Educational. pp.19-25.
- Kvesitadze, E., Sadunishvili, T. & Kvesitadze, G. 2009. Mechanisms of organic contaminants uptake and degradation in plants. *World Academy Science Engineering Technology* 55(6): 458-468.
- Lafabrie, C., Pergent, G., Kantin, R., Pergent-Martini, C. & Gonzalez, J.L. 2007. Trace metals assessment in water, sediment, mussel and seagrass species - Validation of the use of *Posidonia oceanica* as a metal biomonitor. *Chemosphere* 68: 2033-2039.
- Lee, K.S., Park, S.R. & Kim, Y.K. 2007. Effects of irradiance, temperature, and nutrients on growth dynamics of seagrasses: A review. *Journal of Experimental Marine Biology and Ecology* 350: 144-175.
- Lewis, M.A., Dantin, D.D., Chancy, C.A., Abel, K.C. & Lewis, C.G. 2007. Florida seagrass habitat evaluation: A comparative survey for chemical quality. *Environmental Pollution* 146: 206-218.
- Llagostera, I., Pérez, M. & Romero, J. 2011. Trace metal content in the seagrass *Cymodocea nodosa*: Differential accumulation in plant organs. *Aquatic Botany* 95: 124-128.
- Macinnis-Ng, C.M.O. & Ralph, P.J. 2004. Variations in sensitivity to copper and zinc among three isolated populations of the seagrass, *Zostera capricorni*. *Journal of Experimental Marine Biology and Ecology* 302: 63-83.
- Macinnis-Ng, C.M.O. & Ralph, P.J. 2002. Towards a more ecologically relevant assessment of the impact of heavy metals on the photosynthesis of the seagrass, *Zostera capricorni*. *Marine Pollution Bulletin* 45: 100-106.
- Malea, P. 1994. Seasonal variation and local distribution of metals in the seagrass *Halophila stipulacea* (Forsk.) Aschers in the Antikyra Gulf, Greece. *Environmental Pollution* 85: 77-85.
- Malea, P., Haritonidis, S. & Kevrekidis, T. 1994. Seasonal and local variations of metal concentrations in the seagrass *Posidonia oceanica* (L.) Delile in the Antikyra Gulf, Greece. *The Science of the Total Environment* 153: 225-235.
- McKenzie, L.J., Finkbeiner, M.A. & Kirkman, H. 2001. Methods for mapping seagrass distribution. In *Global Seagrass Research Methods*, edited by Short, F.T. & Coles, R.G. Amsterdam: Elsevier Science B.V.
- Nienhuis, P.H. 1986. Background levels of heavy metals in nine tropical seagrass species in Indonesia. *Marine Pollution Bulletin* 17: 508-511.
- Ooi, J.L.S., Kendrick, G.A., Van Niel, K.P. & Affendi, Y.A. 2011. Knowledge gaps in tropical Southeast Asian seagrass systems. *Estuarine, Coastal and Shelf Science* 92: 118-131.
- Prange, J.A. & Dennison, W.C. 2000. Physiological responses of five seagrass species to trace metals. *Marine Pollution Bulletin* 41: 327-336.
- Rensing, C. & Rosen, B.P. 2009. Heavy metals cycle (arsenic, mercury, selenium, others). In *Encyclopedia of Microbiology* (3rd ed), edited by Moselio, S. Oxford: Academic Press.
- Ripperger, S., Rehkämper, M., Porcelli, D. & Halliday, A.N. 2007. Cadmium isotope fractionation in seawater - A signature of biological activity. *Earth and Planetary Science Letters* 261: 670-684.
- Rusli, M.H.B.M. 2012. Protecting vital sea lines of communication: A study of the proposed designation of the Straits of Malacca and Singapore as a particularly sensitive sea area. *Ocean and Coastal Management* 57: 79-94.
- Sanchiz, C.M., García-Carrascosa, A. & Pastor, A. 2000. Heavy metal contents in soft-bottom marine macrophytes and sediments along the Mediterranean Coast of Spain. *Marine Ecology* 21: 1-16.
- Schlacher-Hoenlinger, M.A. & Schlacher, T.A. 1998. Differential accumulation patterns of heavy metals among the dominant macrophytes of a Mediterranean seagrass meadow. *Chemosphere* 37: 1511-1519.
- Short, F.T. & Coles, R.G. 2001. *Global Seagrass Research Methods*. Amsterdam: Elsevier Science B.V. pp. 1-3.
- Short, F.T., Coles, R.G. & Pergent-Martini, C. 2001. Global seagrass distribution. In *Global Seagrass Research Methods*, edited by Short, F.T. & Coles, R.G. Amsterdam: Elsevier Science B.V. pp. 5-30.

- Short, F.T., Polidoro, B., Livingstone, S.R., Carpenter, K.E., Bandeira, S., Bujang, J.S., Calumpong, H.P., Carruthers, T.J.B., Coles, R.G., Dennison, W.C., Erftemeijer, P.L.A., Fortes, M.D., Freeman, A.S., Jagtap, T.G., Kamal, A.H.M., Kendrick, G.A., Judson Kenworthy, W., La Nafie, Y.A., Nasution, I.M., Orth, R.J., Prathep, A., Sanciangco, J.C., Tussenbroek, B.V., Vergara, S.G., Waycott, M. & Zieman, J.C. 2011. Extinction risk assessment of the world's seagrass species. *Biological Conservation* 144: 1961-1971
- Vance, D., Archer, C., Bermin, J., Perkins, J., Statham, P.J., Lohan, M.C., Ellwood, M.J. & Mills, R.A. 2008. The copper isotope geochemistry of rivers and the oceans. *Earth and Planetary Science Letters* 274: 204-213.
- Vermaat, J.E., Rollon, R.N., Lacap, C.D.A., Billot, C., Alberto, F., Nacorda, H.M.E., Wiegman, F. & Terrados, J. 2004. Meadow fragmentation and reproductive output of the SE Asian seagrass *Enhalus acoroides*. *Journal of Sea Research* 52: 321-328.
- Villate, F., Iriarte, A., Uriarte, I., Intxausti, L. & De La Sota, A. 2013. Dissolved oxygen in the rehabilitation phase of an estuary: Influence of sewage pollution abatement and hydro-climatic factors. *Marine Pollution Bulletin* 70: 234-246.
- Waycott, M., Longstaff, B.J. & Mellors, J. 2005. Seagrass population dynamics and water quality in the Great Barrier Reef region: A review and future research directions. *Marine Pollution Bulletin* 51: 343-350.
- Zhan, X-H., Ma, H-L., Zhou, L-X., Liang, J-R. & Xu, G-H. 2010. Accumulation of phenanthrene by roots of intact wheat (*Triticum acstivnm L.*) seedlings: Passive or active uptake?. *BMC Plant Biology* 10: 52.

Department of Environmental Engineering  
Faculty of Civil Engineering  
Universiti Teknologi Malaysia  
81310 Skudai, Johor Darul Takzim  
Malaysia

\*Corresponding author; email: shamila@utm.my

Received: 10 December 2013

Accepted: 5 August 2014